

PUBLICATIONS

- Georges Aad et al. Measurement of the top quark pair production cross-section with ATLAS in the single lepton channel. *Phys.Lett.*, B711:244–263, 2012.
- Georges Aad et al. A measurement of the ratio of the W and Z cross sections with exactly one associated jet in pp collisions at $\sqrt{s} = 7$ TeV with ATLAS. *Phys.Lett.*, B708:221–240, 2012.
- Georges Aad et al. Measurements of underlying-event properties using neutral and charged particles in *pp* collisions at 900 GeV and 7 TeV with the ATLAS detector at the LHC. *Eur.Phys.J.*, C71:1636, 2011.
- Georges Aad et al. Charged-particle multiplicities in pp interactions measured with the ATLAS detector at the LHC. *New J.Phys.*, 13:053033, 2011.
- G. Aad et al. Charged-particle multiplicities in pp interactions at $\sqrt{s} = 900$ GeV measured with the ATLAS detector at the LHC. *Phys.Lett.*, B688:21–42, 2010.
- W. H. Bell et al. Optorsim: A Grid Simulator for Studying Dynamic Data Replication Strategies. *Int. Journal of High Perf. Comp. Apps*, 17:403–416, 2003.
- W. H. Bell et al. Simulation of Dynamic Grid Replication Strategies in OptorSim. *Lecture Notes in Computer Science*, 2536/2002:46–57, 2002.
- K. Anikeev et al. B physics at the Tevatron: Run II and beyond. *FERMILAB-Pub-01/197*, 2001.
- C. Da Via, W. H. Bell, L. Casagrande, V. Granata, and V. G. Palmieri. Cryogenic temperature performance of heavily irradiated silicon detectors. *Nucl. Instrum. Meth.*, A434:114–117, 1999.
- W. H. Bell, L. Casagrande, C. Da Via, V. Granata, and V. G. Palmieri. Temperature dependence of the charge collection efficiency in heavily irradiated silicon detectors. *Nucl. Instrum. Meth.*, A435:187–193, 1999.
- L. Casagrande et al. Tracking with heavily irradiated silicon detectors operated at cryogenic temperatures. *IEEE Trans. Nucl. Sci.*, 46:228–231, 1999.

CO.-AUTHORED

- Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys.Lett.B*, 2012.
- Georges Aad et al. Combined search for the Standard Model Higgs boson in *pp* collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D86:032003, 2012.
- Georges Aad et al. Measurement of the b-hadron production cross section using decays to $D^*\mu^-X$ final states in *pp* collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Nucl.Phys.*, B864:341–381, 2012.
- Georges Aad et al. Hunt for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collisions. *JHEP*, 1207:167, 2012.
- Georges Aad et al. Evidence for the associated production of a W boson and a top quark in ATLAS at $\sqrt{s} = 7$ TeV. *Phys.Lett.B*, 2012.
- Georges Aad et al. A search for $t\bar{t}$ resonances with the ATLAS detector in 2.05 fb^{-1} of proton-proton collisions at $\sqrt{s} = 7$ TeV. *Eur.Phys.J.*, C72:2083, 2012.
- Georges Aad et al. Measurement of the W boson polarization in top quark decays with the ATLAS detector. *JHEP*, 1206:088, 2012.
- Georges Aad et al. Search for $t\bar{b}$ resonances in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.Lett.*, 109:081801, 2012.
- Georges Aad et al. Search for lepton flavour violation in the emu continuum with the ATLAS detector in $\sqrt{s} = 7$ TeV *pp* collisions at the LHC. *Eur.Phys.J.*, C72:2040, 2012.
- Georges Aad et al. Search for scalar top quark pair production in natural gauge mediated supersymmetry models with the ATLAS detector in *pp* collisions at $\sqrt{s} = 7$ TeV. *Phys.Lett.*, B715:44–60, 2012.
- Georges Aad et al. Measurement of τ polarization in $W \rightarrow \tau\nu$ decays with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV. *Eur.Phys.J.*, C72:2062, 2012.
- Georges Aad et al. Search for supersymmetry in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector. *Phys.Rev.Lett.*, 108:261804, 2012.

Georges Aad et al. Search for supersymmetry with jets, missing transverse momentum and at least one hadronically decaying tau lepton in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys. Lett.*, B714:197–214, 2012.

Georges Aad et al. Search for charged Higgs bosons decaying via $H^\pm \rightarrow \tau\nu$ in top quark pair events using pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector. *JHEP*, 1206:039, 2012.

Georges Aad et al. Search for resonant WZ production in the WZ to $l\nu l'$ channel in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector. *Phys. Rev.*, D85:112012, 2012.

Georges Aad et al. Search for pair production of a new quark that decays to a Z boson and a bottom quark with the ATLAS detector. *Phys.Rev.Lett.*, 109:071801, 2012.

Georges Aad et al. Search for the decay $B_s^0 \rightarrow \mu\mu\mu$ with the ATLAS detector. *Phys.Lett.*, B713:387–407, 2012.

Georges Aad et al. Search for events with large missing transverse momentum, jets, and at least two tau leptons in 7 TeV proton-proton collision data with the ATLAS detector. *Phys.Lett.*, B714:180–196, 2012.

Georges Aad et al. Measurement of the WW cross section in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector and limits on anomalous gauge couplings. *Phys.Lett.*, B712:289–308, 2012.

Georges Aad et al. Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b-jets with the ATLAS detector. *Phys.Rev.*, D85:112006, 2012.

Georges Aad et al. Search for gluinos in events with two same-sign leptons, jets and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV. *Phys.Rev.Lett.*, 108:241802, 2012.

Georges Aad et al. Measurement of the top quark mass with the template method in the $t\bar{t} \rightarrow \text{lepton} + \text{jets}$ channel using ATLAS data. *Eur.Phys.J.*, C72:2046, 2012.

Georges Aad et al. Search for heavy neutrinos and right-handed W bosons in events with two leptons and jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Eur.Phys.J.*, C72:2056, 2012.

Georges Aad et al. Measurement of $t\bar{t}$ production with a veto on additional central jet activity in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector. *Eur.Phys.J.*, C72:2043, 2012.

Georges Aad et al. Jet mass and substructure of inclusive jets in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS experiment. *JHEP*, 1205:128, 2012.

Georges Aad et al. Measurement of the charge asymmetry in top quark pair production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector. *Eur.Phys.J.*, C72:2039, 2012.

Georges Aad et al. Observation of spin correlation in $t\bar{t}$ events from pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector. *Phys.Rev.Lett.*, 108:212001, 2012.

Georges Aad et al. Determination of the strange quark density of the proton from ATLAS measurements of the $W \rightarrow l\nu$ and $Z \rightarrow ll$ cross sections. *Phys.Rev.Lett.*, 109:012001, 2012.

Georges Aad et al. Measurement of inclusive two-particle angular correlations in pp collisions with the ATLAS detector at the LHC. *JHEP*, 1205:157, 2012.

Georges Aad et al. Search for second generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. 2012.

Georges Aad et al. Measurement of the production cross section of an isolated photon associated with jets in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:092014, 2012.

Georges Aad et al. Forward-backward correlations and charged-particle azimuthal distributions in pp interactions using the ATLAS detector. *JHEP*, 1207:019, 2012.

Georges Aad et al. Measurement of the azimuthal anisotropy for charged particle production in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector. *Phys.Rev.*, C86:014907, 2012.

Georges Aad et al. Measurement of the polarisation of W bosons produced with large transverse momentum in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment. *Eur.Phys.J.*, C72:2001, 2012.

Georges Aad et al. Search for a light Higgs boson decaying to long-lived weakly-interacting particles in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.Lett.*, 108:251801, 2012.

Georges Aad et al. Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC. 2012.

Georges Aad et al. Search for new particles decaying to ZZ using final states with leptons and jets with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions. *Phys.Lett.*, B712:331–350, 2012.

Georges Aad et al. Search for FCNC single top-quark production at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B712:351–369, 2012.

Georges Aad et al. Measurement of the azimuthal ordering of charged hadrons with the ATLAS detector. 2012.

Georges Aad et al. Search for down-type fourth generation quarks with the ATLAS detector in events with one lepton and hadronically decaying W bosons. *Phys.Rev.Lett.*, 109:032001, 2012.

Georges Aad et al. Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *JHEP*, 1204:069, 2012.

Georges Aad et al. Measurement of the cross section for top-quark pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector using final states with two high-pt leptons. *JHEP*, 1205:059, 2012.

Georges Aad et al. Search for anomaly-mediated supersymmetry breaking with the ATLAS detector based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV. *Eur.Phys.J.*, C72:1993, 2012.

Georges Aad et al. Search for pair-produced heavy quarks decaying to Wq in the two-lepton channel at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D86:012007, 2012.

Georges Aad et al. Search for pair production of a heavy up-type quark decaying to a W boson and a b quark in the lepton+jets channel with the ATLAS detector. *Phys.Rev.Lett.*, 108:261802, 2012.

Georges Aad et al. Search for the Standard Model Higgs boson in the decay channel $H \rightarrow ZZ(*) \rightarrow 4l$ with 4.8 fb⁻¹ of pp collision data at $\sqrt{s} = 7$ TeV with ATLAS. *Phys.Lett.*, B710:383–402, 2012.

Georges Aad et al. Search for the Standard Model Higgs boson in the diphoton decay channel with 4.9 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV with ATLAS. *Phys.Rev.Lett.*, 108:111803, 2012.

Georges Aad et al. Combined search for the Standard Model Higgs boson using up to 4.9 fb⁻¹ of pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC. *Phys.Lett.*, B710:49–66, 2012.

Georges Aad et al. Search for decays of stopped, long-lived particles from 7 TeV pp collisions with the ATLAS detector. *Eur.Phys.J.*, C72:1965, 2012.

Georges Aad et al. Search for excited leptons in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:072003, 2012.

Georges Aad et al. Rapidity gap cross sections measured with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV. *Eur.Phys.J.*, C72:1926, 2012.

Georges Aad et al. Study of jets produced in association with a W boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:092002, 2012.

Georges Aad et al. Search for anomalous production of prompt like-sign muon pairs and constraints on physics beyond the Standard Model with the ATLAS detector. *Phys.Rev.*, D85:032004, 2012.

Georges Aad et al. Jet energy measurement with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV. 2011.

Georges Aad et al. Measurement of inclusive jet and dijet production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector. *Phys. Rev.*, D86:014022, 2012.

Georges Aad et al. Observation of a new χ_b state in radiative transitions to $\Upsilon(1S)$ and $\Upsilon(2S)$ at ATLAS. *Phys. Rev. Lett.*, 108:152001, 2012.

Georges Aad et al. Search for first generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B709:158–176, 2012.

Georges Aad et al. Measurement of $D^{*+/-}$ meson production in jets from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:052005, 2012.

Georges Aad et al. Search for contact interactions in dilepton events from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B712:40–58, 2012.

Georges Aad et al. Search for scalar bottom pair production with the ATLAS detector in pp Collisions at $\sqrt{s} = 7$ TeV. *Phys.Rev.Lett.*, 108:181802, 2012.

Georges Aad et al. Search for production of resonant states in the photon-jet mass distribution using pp collisions at $\sqrt{s} = 7$ TeV collected by the ATLAS detector. *Phys.Rev.Lett.*, 108:211802, 2012.

Georges Aad et al. Search for the Higgs boson in the $H \rightarrow WW(*) \rightarrow l\nu l\nu$ decay channel in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.Lett.*, 108:111802, 2012.

Georges Aad et al. Search for Extra Dimensions using diphoton events in 7 TeV proton-proton collisions with the ATLAS detector. *Phys.Lett.*, B710:538–556, 2012.

Georges Aad et al. Measurement of the WZ production cross section and limits on anomalous triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B709:341–357, 2012.

Georges Aad et al. Search for Diphoton Events with Large Missing Transverse Momentum in 1 fb^{-1} of 7 TeV Proton-Proton Collision Data with the ATLAS Detector. *Phys.Lett.*, B710:519–537, 2012.

Georges Aad et al. Measurement of the production cross section for Z/gamma^* in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:032009, 2012.

Georges Aad et al. Kshort and Lambda production in pp interactions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector at the LHC. *Phys.Rev.*, D85:012001, 2012.

Georges Aad et al. Search for strong gravity signatures in same-sign dimuon final states using the ATLAS detector at the LHC. *Phys.Lett.*, B709:322–340, 2012.

Georges Aad et al. A study of the material in the ATLAS inner detector using secondary hadronic interactions. *JINST*, 7:P01013, 2012.

Georges Aad et al. Searches for supersymmetry with the ATLAS detector using final states with two leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions. *Phys.Lett.*, B709:137–157, 2012.

Georges Aad et al. Measurement of the ZZ production cross section and limits on anomalous neutral triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.Lett.*, 108:041804, 2012.

Georges Aad et al. Electron performance measurements with the ATLAS detector using the 2010 LHC proton-proton collision data. *Eur.Phys.J.*, C72:1909, 2012.

Georges Aad et al. Search for Massive Colored Scalars in Four-Jet Final States in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS Detector. *Eur.Phys.J.*, C71:1828, 2011.

Georges Aad et al. Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector. *JHEP*, 1111:099, 2011.

Georges Aad et al. Performance of the ATLAS Trigger System in 2010. *Eur.Phys.J.*, C72:1849, 2012.

Georges Aad et al. Measurement of the inclusive and dijet cross-sections of b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Eur.Phys.J.*, C71:1846, 2011.

Georges Aad et al. Search for supersymmetry in final states with jets, missing transverse momentum and one isolated lepton in $\sqrt{s} = 7$ TeV pp collisions using 1 fb^{-1} of ATLAS data. *Phys.Rev.*, D85:012006, 2012.

Georges Aad et al. Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions. *Phys.Lett.*, B710:67–85, 2012.

Georges Aad et al. Search for the Standard Model Higgs boson in the decay channel $H \rightarrow ZZ(*) \rightarrow 4l$ with the ATLAS detector. *Phys.Lett.*, B705:435–451, 2011.

Georges Aad et al. Measurement of the jet fragmentation function and transverse profile in proton-proton collisions at a center-of-mass energy of 7 TeV with the ATLAS detector. *Eur.Phys.J.*, C71:1795, 2011.

Georges Aad et al. Measurement of the inclusive W^\pm and Z/γ cross sections in the electron and muon decay channels in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:072004, 2012.

Georges Aad et al. Search for New Phenomena in $t\bar{t}$ Events With Large Missing Transverse Momentum in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector. *Phys.Rev.Lett.*, 108:041805, 2012.

Georges Aad et al. Search for the Higgs boson in the $H \rightarrow WW \rightarrow lvjj$ decay channel in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.Lett.*, 107:231801, 2011.

Georges Aad et al. Search for a Standard Model Higgs boson in the $H \rightarrow ZZ \rightarrow l^+l^-v\nu^-$ decay channel with the ATLAS detector. *Phys.Rev.Lett.*, 107:221802, 2011.

Georges Aad et al. Search for a heavy neutral particle decaying into an electron and a muon using 1 fb⁻¹ of ATLAS data. *Eur.Phys.J.*, C71:1809, 2011.

Georges Aad et al. Search for displaced vertices arising from decays of new heavy particles in 7 TeV pp collisions at ATLAS. *Phys.Lett.*, B707:478–496, 2012.

Georges Aad et al. Measurement of the cross-section for b-jets produced in association with a Z boson at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B706:295–313, 2012.

Georges Aad et al. Measurement of the cross section for the production of a W boson in association with b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B707:418–437, 2012.

Georges Aad et al. Measurements of the electron and muon inclusive cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B707:438–458, 2012.

Georges Aad et al. Search for New Physics in the Dijet Mass Distribution using 1 fb⁻¹ of pp Collision Data at $\sqrt{s} = 7$ TeV collected by the ATLAS Detector. *Phys.Lett.*, B708:37–54, 2012.

Georges Aad et al. Measurement of the Transverse Momentum Distribution of W Bosons in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector. *Phys.Rev.*, D85:012005, 2012.

Georges Aad et al. Measurement of the centrality dependence of the charged particle pseudorapidity distribution in lead-lead collisions at $\sqrt{\langle s_{NN} \rangle} = 2.76$ TeV with the ATLAS detector. *Phys.Lett.*, B710:363–382, 2012.

Georges Aad et al. Measurement of the pseudorapidity and transverse momentum dependence of the elliptic flow of charged particles in lead-lead collisions at $\sqrt{\langle s_{NN} \rangle} = 2.76$ TeV with the ATLAS detector. *Phys.Lett.*, B707:330–348, 2012.

Georges Aad et al. Search for the Standard Model Higgs boson in the two photon decay channel with the ATLAS detector at the LHC. *Phys.Lett.*, B705:452–470, 2011.

Georges Aad et al. Performance of Missing Transverse Momentum Reconstruction in Proton-Proton Collisions at 7 TeV with ATLAS. *Eur.Phys.J.*, C72:1844, 2012.

Georges Aad et al. Search for a heavy Standard Model Higgs boson in the channel $H \rightarrow ZZ \rightarrow llqq$ using the ATLAS detector. *Phys.Lett.*, B707:27–45, 2012.

Georges Aad et al. Measurement of the W to tau nu Cross Section in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment. *Phys.Lett.*, B706:276–294, 2012.

Georges Aad et al. Measurement of the top quark pair production cross section in pp collisions at $\sqrt{s} = 7$ TeV in dilepton final states with ATLAS. *Phys.Lett.*, B707:459–477, 2012.

Georges Aad et al. Measurement of the Z to tau tau Cross Section with the ATLAS Detector. *Phys.Rev.*, D84:112006, 2011.

Georges Aad et al. Search for dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.Lett.*, 107:272002, 2011.

Georges Aad et al. Search for a heavy gauge boson decaying to a charged lepton and a neutrino in 1 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector. *Phys.Lett.*, B705:28–46, 2011.

Georges Aad et al. Inclusive search for same-sign dilepton signatures in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *JHEP*, 1110:107, 2011.

Georges Aad et al. Measurement of the inclusive isolated prompt photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV using 35 pb^{-1} of ATLAS data. *Phys.Lett.*, B706:150–167, 2011.

Georges Aad et al. Search for neutral MSSM Higgs bosons decaying to $\tau^+\tau^-$ pairs in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B705:174–192, 2011.

Georges Aad et al. Properties of jets measured from tracks in proton-proton collisions at center-of-mass energy $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D84:054001, 2011.

Georges Aad et al. Measurement of the transverse momentum distribution of Z/gamma* bosons in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B705:415–434, 2011.

Georges Aad et al. Measurement of multi-jet cross sections in proton-proton collisions at a 7 TeV center-of-mass energy. *Eur.Phys.J.*, C71:1763, 2011.

Georges Aad et al. Measurement of dijet production with a veto on additional central jet activity in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector. *JHEP*, 1109:053, 2011.

Georges Aad et al. Search for Diphoton Events with Large Missing Transverse Energy with 36 pb^{-1} of 7 TeV Proton-Proton Collision Data with the ATLAS Detector. *Eur.Phys.J.*, C71:1744, 2011.

Georges Aad et al. Measurement of the isolated di-photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D85:012003, 2012.

Georges Aad et al. Measurement of the Upsilon(1S) Production Cross-Section in pp Collisions at $\sqrt{s} = 7$ TeV in ATLAS. *Phys.Lett.*, B705:9–27, 2011.

Georges Aad et al. Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions. *Phys.Lett.*, B705:294–312, 2011.

Georges Aad et al. Search for Heavy Long-Lived Charged Particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV. *Phys.Lett.*, B703:428–446, 2011.

Georges Aad et al. Measurement of the differential cross-sections of inclusive, prompt and non-prompt J/psi production in proton-proton collisions at $\sqrt{s} = 7$ TeV. *Nucl.Phys.*, B850:387–444, 2011.

Georges Aad et al. Search for an excess of events with an identical flavour lepton pair and significant missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS detector. *Eur.Phys.J.*, C71:1647, 2011.

Georges Aad et al. Search for high mass dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment. *Phys.Lett.*, B700:163–180, 2011.

Georges Aad et al. Search for New Physics in Dijet Mass and Angular Distributions in pp Collisions at $\sqrt{s} = 7$ TeV Measured with the ATLAS Detector. *New J.Phys.*, 13:053044, 2011.

Georges Aad et al. Search for heavy vector-like quarks coupling to light quarks in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B712:22–39, 2012.

Georges Aad et al. Search for stable hadronising squarks and gluinos with the ATLAS experiment at the LHC. *Phys.Lett.*, B701:1–19, 2011.

Georges Aad et al. Search for high-mass states with one lepton plus missing transverse momentum in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B701:50–69, 2011.

Georges Aad et al. Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions. *Phys.Lett.*, B701:186–203, 2011.

Georges Aad et al. Measurement of Dijet Azimuthal Decorrelations in pp Collisions at $\sqrt{s} = 7$ TeV. *Phys. Rev. Lett.*, 106:172002, 2011.

Georges Aad et al. Search for supersymmetry using final states with one lepton, jets, and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV pp. *Phys.Rev.Lett.*, 106:131802, 2011.

Georges Aad et al. Search for Massive Long-lived Highly Ionising Particles with the ATLAS Detector at the LHC. *Phys.Lett.*, B698:353–370, 2011.

Georges Aad et al. Luminosity Determination in pp Collisions at $\sqrt{s} = 7$ TeV Using the ATLAS Detector at the LHC. *Eur.Phys.J.*, C71:1630, 2011.

Georges Aad et al. Study of Jet Shapes in Inclusive Jet Production in pp Collisions at $\sqrt{s} = 7$ TeV using the ATLAS Detector. *Phys.Rev.*, D83:052003, 2011.

Georges Aad et al. Measurement of the production cross section for W-bosons in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Lett.*, B698:325–345, 2011.

Georges Aad et al. Measurement of the centrality dependence of J/ψ yields and observation of Z production in lead-lead collisions with the ATLAS detector at the LHC. *Phys.Lett.*, B697:294–312, 2011.

Georges Aad et al. Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Phys.Rev.*, D83:052005, 2011.

Georges Aad et al. Search for Diphoton Events with Large Missing Transverse Energy in 7 TeV Proton-Proton Collisions with the ATLAS Detector. *Phys.Rev.Lett.*, 106:121803, 2011.

Georges Aad et al. Measurement of the top quark-pair production cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV. *Eur.Phys.J.*, C71:1577, 2011.

Georges Aad et al. Measurement of underlying event characteristics using charged particles in pp collisions at $\sqrt{s} = 900$ GeV and 7 TeV with the ATLAS detector. *Phys. Rev. D*, 83:112001, 2011.

Georges Aad et al. Observation of a Centrality-Dependent Dijet Asymmetry in Lead-Lead Collisions at $\sqrt{S(NN)} = 2.76$ TeV with the ATLAS Detector at the LHC. *Phys.Rev.Lett.*, 105:252303, 2010.

Georges Aad et al. Measurement of the $W \rightarrow l\nu$ and $Z/\gamma^* \rightarrow ll$ production cross sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *JHEP*, 1012:060, 2010.

Georges Aad et al. Search for Quark Contact Interactions in Dijet Angular Distributions in pp Collisions at $\sqrt{s} = 7$ TeV Measured with the ATLAS Detector. *Phys.Lett.*, B694:327–345, 2011.

Georges Aad et al. Measurement of inclusive jet and dijet cross sections in proton-proton collisions at 7 TeV centre-of-mass energy with the ATLAS detector. *Eur.Phys.J.*, C71:1512, 2011.

Georges Aad et al. Search for New Particles in Two-Jet Final States in 7 TeV Proton-Proton Collisions with the ATLAS Detector at the LHC. *Phys.Rev.Lett.*, 105:161801, 2010.

G Aad et al. Performance of the ATLAS Detector using First Collision Data. *JHEP*, 1009:056, 2010.

G. Aad et al. Drift Time Measurement in the ATLAS Liquid Argon Electromagnetic Calorimeter using Cosmic Muons. *Eur.Phys.J.*, C70:755–785, 2010.

G. Aad et al. Readiness of the ATLAS Liquid Argon Calorimeter for LHC Collisions. *Eur.Phys.J.*, C70:723–753, 2010.

P. Ryan et al. The ATLAS Inner Detector commissioning and calibration. *Eur. Phys. J.*, C70:787–821, 2010.

A. Abdesselam et al. The integration and engineering of the ATLAS Semiconductor Tracker barrel. *JINST*, 3:P10006, 2008.

- E. Abat et al. Combined performance tests before installation of the ATLAS Semiconductor and Transition Radiation Tracking Detectors. *JINST*, 3:P08003, 2008.
- A. Abdesselam et al. Engineering for the ATLAS SemiConductor Tracker (SCT) end-cap. *JINST*, 3:P05002, 2008.
- G. Aad et al. The ATLAS Experiment at the CERN Large Hadron Collider. *JINST*, 3:S08003, 2008.
- M. Abolins et al. Integration of the trigger and data acquisition systems in ATLAS. *J. Phys. Conf. Ser.*, 119:022001, 2008.
- T. Fonseca-Martin et al. Event reconstruction algorithms for the ATLAS trigger. *J. Phys. Conf. Ser.*, 119:022022, 2008.
- Maris A. Abolins et al. The ATLAS trigger: Commissioning with cosmic rays. *J. Phys. Conf. Ser.*, 119:022014, 2008.
- P. J. W. Faulkner et al. GridPP: Development of the UK computing Grid for particle physics. *J. Phys.*, G32:N1–N20, 2006.
- Darin E. Acosta et al. Search for W and Z bosons in the reaction $\bar{p}p \rightarrow 2\text{jets} + \gamma$ at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D73:012001, 2006.
- Darin E. Acosta et al. A search for supersymmetric Higgs bosons in the di-tau decay mode in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D72:072004, 2005.
- Darin E. Acosta et al. K_S^0 and Λ^0 production studies in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ -GeV and 630-GeV. *Phys. Rev.*, D72:052001, 2005.
- Darin E. Acosta et al. Measurement of the azimuthal angle distribution of leptons from W boson decays as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D73:052002, 2006.
- Darin E. Acosta et al. Search for Higgs bosons decaying into $b\bar{b}$ and produced in association with a vector boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 95:051801, 2005.
- Darin E. Acosta et al. Measurements of $b\bar{b}$ azimuthal production correlations in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D71:092001, 2005.
- Darin E. Acosta et al. Measurement of the W boson polarization in top decay at CDF at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D71:031101, 2005.
- Darin E. Acosta et al. Measurement of charged particle multiplicities in gluon and quark jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 94:171802, 2005.
- Darin E. Acosta et al. Comparison of three-jet events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV to predictions from a next-to-leading order QCD calculation. *Phys. Rev.*, D71:032002, 2005.
- Darin E. Acosta et al. Inclusive search for anomalous production of high p_T like-sign lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 93:061802, 2004.
- Darin E. Acosta et al. Direct photon cross section with conversions at CDF. *Phys. Rev.*, D70:074008, 2004.
- Darin E. Acosta et al. The underlying event in hard interactions at the Tevatron $\bar{p}p$ collider. *Phys. Rev.*, D70:072002, 2004.
- Darin E. Acosta et al. Optimized search for single top quark production at the Fermilab Tevatron. *Phys. Rev.*, D69:052003, 2004.
- Darin E. Acosta et al. Observation of the narrow state $X(3872) \rightarrow J/\psi\pi^+\pi^-$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV. *Phys. Rev. Lett.*, 93:072001, 2004.
- D. Acosta et al. Heavy flavor properties of jets produced in $p\bar{p}$ interactions at $\sqrt{s} = 1.8$ -TeV. *Phys. Rev.*, D69:072004, 2004.
- Darin E. Acosta et al. Measurement of the polar-angle distribution of leptons from W boson decay as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D70:032004, 2004.
- V. M. Abazov et al. Combination of CDF and DØ results on W boson mass and width. *Phys. Rev.*, D70:092008, 2004.
- Darin E. Acosta et al. Inclusive double pomeron exchange at the Fermilab Tevatron $\bar{p}p$ collider. *Phys. Rev. Lett.*, 93:141601, 2004.

M. Abreu et al. Recent progress in low-temperature silicon detectors. *Nucl.Phys.Proc.Suppl.*, 125:169–174, 2003.

V. G. Palmieri et al. Cryogenic semiconductor high-intensity radiation monitors. *Nucl. Instrum. Meth.*, A510:97–100, 2003.

D. Acosta et al. Measurement of the average time-integrated mixing probability of b -flavored hadrons produced at the Tevatron. *Phys. Rev.*, D69:012002, 2004.

Darin E. Acosta et al. Search for the flavor-changing neutral current decay $D^0 \rightarrow \mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. *Phys. Rev.*, D68:091101, 2003.

Darin E. Acosta et al. Measurement of the mass difference $m(D_s^+) - m(D^+)$ at CDF II. *Phys. Rev.*, D68:072004, 2003.

Darin E. Acosta et al. Measurement of prompt charm meson production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. *Phys. Rev. Lett.*, 91:241804, 2003.

Darin E. Acosta et al. Search for associated production of Υ and vector boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 90:221803, 2003.

Darin E. Acosta et al. Search for lepton flavor violating decays of a heavy neutral particle in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 91:171602, 2003.

Darin E. Acosta et al. Search for pair production of scalar top quarks in R -parity violating decay modes in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 92:051803, 2004.

Darin E. Acosta et al. Central pseudorapidity gaps in events with a leading antiproton at the Fermilab Tevatron $p\bar{p}$ collider. *Phys. Rev. Lett.*, 91:011802, 2003.

Darin E. Acosta et al. Search for the supersymmetric partner of the top quark in dilepton events from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 90:251801, 2003.

Darin E. Acosta et al. Cross section for forward J/ψ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D66:092001, 2002.

Darin E. Acosta et al. Search for long-lived charged massive particles in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 90:131801, 2003.

T. O. Niinikoski et al. Radiation hardness of cryogenic silicon detectors. *Nucl. Instrum. Meth.*, A476:569–582, 2002.

G. Ruggiero et al. Silicon detectors irradiated 'in situ' at cryogenic temperatures. *Nucl. Instrum. Meth.*, A476:583–587, 2002.

Darin E. Acosta et al. Search for a W' boson decaying to a top and bottom quark pair in 1.8 TeV $p\bar{p}$ collisions. *Phys. Rev. Lett.*, 90:081802, 2003.

Darin E. Acosta et al. Search for radiative b -hadron decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D66:112002, 2002.

Darin E. Acosta et al. Momentum distribution of charged particles in jets in dijet events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV and comparisons to perturbative QCD predictions. *Phys. Rev.*, D68:012003, 2003.

Darin E. Acosta et al. Measurement of the ratio of b quark production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 630$ GeV and $\sqrt{s} = 1800$ GeV. *Phys. Rev.*, D66:032002, 2002.

Darin E. Acosta et al. Branching ratio measurements of exclusive B^+ decays to charmonium with the Collider Detector at Fermilab. *Phys. Rev.*, D66:052005, 2002.

Darin E. Acosta et al. Limits on extra dimensions and new particle production in the exclusive photon and missing energy signature in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 89:281801, 2002.

Darin E. Acosta et al. Measurement of B meson lifetimes using fully reconstructed B decays produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D65:092009, 2002.

L. Casagrande et al. Radiation hard cryogenic silicon detectors. *Nucl. Instrum. Meth.*, A477:299–303, 2002.

Darin E. Acosta et al. Search for new physics in photon lepton events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV. *Phys. Rev. Lett.*, 89:041802, 2002.

Darin E. Acosta et al. Upsilon production and polarization in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV. *Phys. Rev. Lett.*, 88:161802, 2002.

Darin E. Acosta et al. Comparison of the isolated direct photon cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV and $\sqrt{s} = 0.63$ -TeV. *Phys. Rev.*, D65:112003, 2002.

Darin E. Acosta et al. Soft and hard interactions in $p\bar{p}$ collisions at $\sqrt{s} = 1800\text{-GeV}$ and 630-GeV . *Phys. Rev.*, D65:072005, 2002.

Darin E. Acosta et al. Measurement of the B^+ total cross section and B^+ differential cross section $d\sigma/dp_T$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{-TeV}$. *Phys. Rev.*, D65:052005, 2002.

Darin E. Acosta et al. Search for single top quark production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{-TeV}$. *Phys. Rev.*, D65:091102, 2002.

Darin E. Acosta et al. Search for the decay $B_s \rightarrow \mu^+ \mu^- \phi$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{-TeV}$. *Phys. Rev.*, D65:111101, 2002.

Anthony Allen Affolder et al. Charged jet evolution and the underlying event in $p\bar{p}$ collisions at 1.8 TeV . *Phys. Rev.*, D65:092002, 2002.

Darin E. Acosta et al. Search for new physics in photon lepton events in proton antiproton collisions at $\sqrt{s} = 1.8\text{-TeV}$. *Phys. Rev.*, D66:012004, 2002.

Anthony Allen Affolder et al. Diffractive dijet production at $\sqrt{s} = 630\text{ GeV}$ and 1800 GeV at the Fermilab Tevatron. *Phys. Rev. Lett.*, 88:151802, 2002.

D. Acosta et al. Study of the heavy flavor content of jets produced in association with W bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev.*, D65:052007, 2002.

Anthony Allen Affolder et al. Measurement of the strong coupling constant from inclusive jet production at the Tevatron $p\bar{p}$ collider. *Phys. Rev. Lett.*, 88:042001, 2002.

Anthony Allen Affolder et al. A study of $B^0 \rightarrow J/\psi K^{(*)0} \pi^+ \pi^-$ decays with the Collider Detector at Fermilab. *Phys. Rev. Lett.*, 88:071801, 2002.

Anthony Allen Affolder et al. Search for new heavy particles in the WZ^0 final state in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{TeV}$. *Phys. Rev. Lett.*, 88:071806, 2002.

Anthony Allen Affolder et al. Observation of diffractive J/ψ production at the Fermilab Tevatron. *Phys. Rev. Lett.*, 87:241802, 2001.

Anthony Allen Affolder et al. Charged particle multiplicity in jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev. Lett.*, 87:211804, 2001.

Anthony Allen Affolder et al. Search for quark lepton compositeness and a heavy W' boson using the $e\nu$ channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev. Lett.*, 87:231803, 2001.

Anthony Allen Affolder et al. Searches for new physics in events with a photon and b- quark jet at CDF. *Phys. Rev.*, D65:052006, 2002.

L. Casagrande et al. Review on the development of cryogenic silicon detectors. *Nucl. Instrum. Meth.*, A461:150–154, 2001.

Anthony Allen Affolder et al. Search for gluinos and squarks using like-sign dileptons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev. Lett.*, 87:251803, 2001.

Anthony Allen Affolder et al. Measurement of $d(\sigma)/dM$ and forward-backward charge asymmetry for high mass Drell-Yan e^+e^- pairs from $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev. Lett.*, 87:131802, 2001.

Anthony Allen Affolder et al. Cross section and heavy quark composition of $\gamma + \mu$ events produced in $p\bar{p}$ collisions. *Phys. Rev.*, D65:012003, 2002.

Anthony Allen Affolder et al. Search for gluinos and scalar quarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$ using the missing energy plus multijets signature. *Phys. Rev. Lett.*, 88:041801, 2002.

Anthony Allen Affolder et al. Production of χ_{c1} and χ_{c2} in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev. Lett.*, 86:3963–3968, 2001.

Anthony Allen Affolder et al. Double diffraction dissociation at the Fermilab Tevatron collider. *Phys. Rev. Lett.*, 87:141802, 2001.

Anthony Allen Affolder et al. Search for narrow diphoton resonances and for $\gamma\gamma + W/Z$ signatures in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev.*, D64:092002, 2001.

Anthony Allen Affolder et al. Measurement of the inclusive jet cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev.*, D64:032001, 2001.

Anthony Allen Affolder et al. Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$. *Phys. Rev.*, D64:032002, 2001.

- V. Granata et al. Cryogenic technology for tracking detectors. *Nucl. Instrum. Meth.*, A461:197–199, 2001.
- K. Borer et al. Charge collection efficiency of irradiated cryogenic double-p silicon detector. *Nucl. Instrum. Meth.*, A462:474–483, 2001.
- P. Collins et al. Cryogenic operation of silicon detectors. *Nucl. Instrum. Meth.*, A447:151–159, 2000.
- Anthony Allen Affolder et al. First measurement of the ratio $B(t \rightarrow Wb)/B(t \rightarrow Wq)$ and associated limit on the CKM element $|V_{tb}|$. *Phys. Rev. Lett.*, 86:3233–3238, 2001.
- Anthony Allen Affolder et al. Measurement of the two-jet differential cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ GeV. *Phys. Rev.*, D64:012001, 2001.
- Anthony Allen Affolder et al. Search for the supersymmetric partner of the top quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev.*, D63:091101, 2001.
- Anthony Allen Affolder et al. Search for neutral supersymmetric Higgs bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. *Phys. Rev. Lett.*, 86:4472–4478, 2001.
- Anthony Allen Affolder et al. Test of enhanced leading order QCD in W boson plus jets events from 1.8 TeV $p\bar{p}$ collisions. *Phys. Rev.*, D63:072003, 2001.
- K. Borer, S. Janos, V.G. Palmieri, J. Buytaert, V. Chabaud, et al. Charge collection efficiency and resolution of an irradiated double sided silicon microstrip detector operated at cryogenic temperatures. *Nucl.Instrum.Meth.*, A440:17–37, 2000.
- K. Borer et al. Charge collection efficiency of irradiated silicon detectors operated at cryogenic temperatures. *Nucl. Instrum. Meth.*, A440:5–16, 2000.